This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):


- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

FIGURE 1






Glucose in db/db and ob/ob Diabetic Mice No Treatment ▼62% glucose 5 10 15 20 25 **Treatment Days** ob/ob Mice Blood Glucose, mg/dl - Treatment -O- Vehicle Oral dose 50 mg/kg n = 6 ▼ No treatment ▼ 52% Glucose 5 10 15 20 25 30 Treatment Days db/db Mice 150-100+ 400 Blood Glucose, mg/dl

FIG. 3A

FIG. 3B



Fime Course






FIGURE 5



Serum AST/GOT and ALT/GPT in ob/o b Mice









FIGURE 8





--- Compound VIII + (0.1 µg/ml) LPS ->- Rosiglitazone + (0.1 µg/ml) LPS







--- Rosiglitazone + (0.1 µg/ml) LPS









Normal	Stiff tail	Limp tail	Limp tail with inability to right	Paralysis of one limb	Paralysis of one limb and weakness of 2nd limb	Paralysis of both hind limbs	Moribund	Death
0	0.5	1.0	1.5	2.0	2.5	3.0	4.0	5.0
	0 Normal							